## **AMENDMENTS TO THE CLAIMS:**

The following Listing of Claims replaces all prior Listings and versions of claims in the above-identified application.

## **Listing of Claims**

- 1. (Currently Amended) A method to produce <u>N-acetylglucosamine-6-</u> <u>phosphate glucosamine</u> or N-acetylglucosamine by fermentation, comprising:
  - a) culturing in a fermentation medium a <u>microorganism that is</u> transformed with at least one recombinant nucleic acid molecule comprising a <u>nucleic acid sequence encoding a glucosamine-6-phosphate acetyltransferase</u> that has an amino acid sequence that is at least 95% identical to SEQ ID NO:30 bacterium or yeast which comprises at least one genetic modification that results in the increased expression of a bacterial or yeast glucosamine-6-phosphate acetyltransferase; and
  - b) collecting a product produced from the step of culturing which is selected from the group consisting of <del>glucosamine-6-phosphate, glucosamine, glucosamine-1-phosphate, N-acetylglucosamine-1-phosphate, N-acetylglucosamine-1-phosphate, N-acetylglucosamine-6-phosphate[[,]] and N-acetylglucosamine.</del>
  - 2-3. (Canceled).
- 4. (Currently Amended) The method of Claim 1 [[3]], wherein the recombinant nucleic acid molecule further comprises a non-native promoter.
  - 5-7. (Canceled).
- 8. (Currently Amended) The method of Claim 1 [[3]], wherein the glucosamine-6-phosphate acetyltransferase has the amino acid sequence selected from the group consisting of SEQ ID NO:30., SEQ ID NO:32 and SEQ ID NO:34.

- 9. (Currently Amended) The method of Claim 1 [[3]], wherein expression of the recombinant nucleic acid molecule is inducible.
- 10. (Previously Presented) The method of Claim 9, wherein expression of the recombinant nucleic acid molecule is inducible by lactose.
  - 11. (Canceled).
- 12. (Currently Amended) The method of Claim 10 [[11]], wherein the microorganism genetic modification further comprises a partial or complete deletion or inactivation of a gene encoding a Lacl repressor protein.
  - 13-20. (Canceled).
- 21. (Currently Amended) The method of Claim 1, wherein the <u>microorganism</u> bacterium or yeast further comprises at least one <u>a</u> partial or complete deletion of an endogenous gene encoding a glucosamine-6-phosphate deaminase in the bacterium or yeast that decreases the activity of glucosamine-6-phosphate deaminase in the bacterium or yeast.
  - 22-24. (Canceled).
- 25. (Previously Presented) The method of Claim 1, wherein the step of culturing includes the step of maintaining the carbon source at a concentration of from about 0.5% to about 5% in the fermentation medium.
- 26. (Previously Presented) The method of Claim 1, wherein the step of culturing is performed in a fermentation medium comprising yeast extract.
  - 27. (Previously Presented) The method of Claim 1, wherein the step of

culturing is performed in a fermentation medium comprising a carbon source selected from the group consisting of glucose, fructose, a pentose sugar, lactose and gluconic acid.

- 28. (Previously Presented) The method of Claim 27, wherein the pentose sugar is selected from the group consisting of ribose, xylose, and arabinose.
- 29. (Previously Presented) The method of Claim 1, wherein the step of culturing is performed in a fermentation medium comprising glucose and ribose.
- 30. (Previously Presented) The method of Claim 1, wherein the step of culturing is performed in a fermentation medium comprising glucose and gluconic acid.
- 31. (Previously Presented) The method of Claim 1, wherein the step of culturing is performed at a temperature of from about 25 ℃ to about 45 ℃.
- 32. (Previously Presented) The method of Claim 1, wherein the step of culturing is performed at about 37°C.
- 33. (Previously Presented) The method of Claim 1, wherein the step of culturing is performed at a pH of from about pH 4 to about pH 7.5.
- 34. (Previously Presented) The method of Claim 1, wherein the step of culturing is performed at a pH of from about pH 6.7 to about pH 7.5.
- 35. (Previously Presented) The method of Claim 1, wherein the step of culturing is performed at a pH of from about pH 4.5 to about pH 5.
- 36. (Currently Amended) The method of Claim 1, wherein the <u>microorganism</u> is a bacterium or yeast is a bacterium.

- 37. (Currently Amended) The method of Claim 1, wherein the <u>microorganism</u> is a bacterium or yeast is a yeast.
- 38. (Currently Amended) The method of Claim <u>37</u> [[36]], wherein the bacterium is a bacterium from a genus selected from the group consisting of: *Escherichia, Bacillus, Lactobacillus, Pseudomonas* and *Streptomyces*.
- 39. (Currently Amended) The method of Claim <u>37</u> [[36]], wherein the bacterium is a bacterium from a species selected from the group consisting of: *Escherichia coli*, *Bacillus subtilis*, *Bacillus licheniformis*, *Lactobacillus brevis*, *Pseudomonas aeruginosa* and *Streptomyces lividans*.
- 40. (Currently Amended) The method of Claim <u>251</u> [[37]], wherein the yeast is a yeast from a genus selected from the group consisting of: *Saccharomyces*, *Candida*, *Hansenula*, *Pichia*, *Kluveromyces*, and *Phaffia*.
- 41. (Currently Amended) The method of Claim <u>251</u> [[37]], wherein the yeast is a yeast from a species selected from the group consisting of: *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Candida albicans*, *Hansenula polymorpha*, *Pichia pastoris*, *P. canadensis*, *Kluyveromyces marxianus* and *Phaffia rhodozyma*.

42-44. (Canceled).

- 45. (Currently Amended) The method of Claim 1, wherein the <u>microorganism</u> bacterium or yeast is transformed with a recombinant nucleic acid molecule comprising a nucleic acid sequence encoding a bacterial or yeast phosphoglucoisomerase.
- 46. (Currently Amended) The method of Claim 45, wherein the phosphoglucoisomerase comprises the amino acid sequence of SEQ ID NO:105.

- 47. (Currently Amended) The method of Claim 1, wherein the <u>microorganism</u> bacterium or yeast further comprises a partial or complete deletion of <u>an endogenous</u> gene encoding a phosphofructokinase in the bacterium or yeast.
  - 48. (Canceled).
- 49. (Currently Amended) The method of Claim 1, wherein the <u>microorganism</u> bacterium or yeast has been transformed with a recombinant nucleic acid molecule comprising a nucleic acid sequence encoding a bacterial or yeast glutamine synthetase.
- 50. (Previously Presented) The method of Claim 49, wherein the glutamine synthetase comprises the amino acid sequence of SEQ ID NO:89.
  - 51. (Canceled).
- 52. (Currently Amended) The method of Claim 1, wherein the <u>microorganism</u> bacterium or yeast has been transformed with a recombinant nucleic acid molecule comprising a nucleic acid sequence encoding <u>a bacterial or yeast</u> glucose-6-phosphate dehydrogenase.
- 53. (Previously Presented) The method of Claim 52, wherein the glucose-6-phosphate dehydrogenase comprises the amino acid sequence of SEQ ID NO:95.
- 54. (Currently Amended) The method of Claim 1, wherein the <u>microorganism</u> bacterium or yeast further comprises a partial or complete deletion of at least one <u>endogenous</u> gene encoding an enzyme involved in glycogen synthesis <u>selected from the group consisting of: ADP-glucose pyrophosphorylase, glycogen synthase and a branching enzyme in the bacterium or yeast.</u>

## 55-56. (Canceled).

- 57. (Currently Amended) The method of Claim 1, <u>further comprising</u> wherein the step of collecting comprises recovering an intracellular product from the <u>microorganism</u> bacterium or yeast selected from the group consisting of: intracellular glucosamine-6-phosphate, glucosamine-1-phosphate, N-acetylglucosamine-6-phosphate, N-acetylglucosamine and glucosamine. or recovering an extracellular product from the fermentation medium selected from the group consisting of: glucosamine and N-acetylglucosamine.
- 58. (Currently Amended) The method of Claim 1, further comprising a step selected from the group consisting of:
  - a) purifying a product selected from the group consisting of glucosamine and N-acetylglucosamine from the fermentation medium;
  - b) dephosphorylating a product selected from the group consisting of glucosamine-6-phosphate and glucosamine-1-phosphate to produce glucosamine;
  - e) b) dephosphorylating a product selected from the group consisting of N-acetylglucosamine-6-phosphate and N-acetylglucosamine-1-phosphate to produce N-acetylglucosamine; and
  - d) c) treating a product selected from the group consisting of N-acetylglucosamine, N-acetylglucosamine-6-phosphate and N-acetylglucosamine-1-phosphate to produce a glucosamine product selected from the group consisting of: glucosamine[[,]] and glucosamine HCI. glucosamine-6-phosphate and glucosamine-1-phosphate.
- 59. (Currently Amended) The method of Claim 58, wherein step (c) (d) comprises hydrolyzing the product selected from the group consisting of N-acetylglucosamine[[,]] and N-acetylglucosamine-6-phosphate and N-acetylglucosamine-1-phosphate, under acid and heat conditions or by enzymatic deacetylation.

- 60. (Previously Presented) The method of Claim 1, wherein N-acetylglucosamine produced by the fermentation method is recovered by precipitating N-acetylglucosamine-containing solids from the fermentation broth.
- 61. (Previously Presented) The method of Claim 1, wherein N-acetylglucosamine produced by the fermentation method is recovered by crystallizing N-acetylglucosamine-containing solids from the fermentation broth.

62-218. (Canceled).

- 219. (Currently Amended) A method to produce <u>N-acetylglucosamine-6-</u> <u>phosphate glucosamine</u> or N-acetylglucosamine by fermentation, comprising:
  - a) culturing in a fermentation medium a <u>microorganism</u> <del>bacterium or yeast</del> that expresses:
    - i) a recombinant nucleic acid molecule encoding a glucosamine-6-phosphate acetyltransferase that has an amino acid sequence that is at least 95% identical to SEQ ID NO:30 and has glucosamine-6-phosphate acetyltransferase enzymatic activity; and
    - ii) a recombinant nucleic acid molecule encoding a glucosamine-6-phosphate synthase that has an amino acid sequence that is at least 95% identical to the amino acid sequence selected from the group consisting of: SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, and SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, and SEQ ID NO:20; and has glucosamine-6-phosphate synthase enzymatic activity; and
  - b) collecting a product produced from the step of culturing which is selected from the group consisting of glucosamine-6-phosphate, glucosamine, glucosamine-1-phosphate, N-acetylglucosamine-1-phosphate, N-acetylglucosamine-6-phosphate[[,]] and N-acetylglucosamine.

- 220. (Previously Presented) The method of Claim 219, wherein the glucosamine-6-phosphate acetyltransferase has the amino acid sequence of SEQ ID NO:30.
- 221. (Previously Presented) The method of Claim 219, wherein the glucosamine-6-phosphate synthase has the amino acid sequence of SEQ ID NO:6.
- 222. (Currently Amended) A method to produce <u>N-acetylglucosamine-6-phosphate glucosamine</u> or N-acetylglucosamine by fermentation, comprising:
  - a) culturing in a fermentation medium an *E. coli* that expresses:
  - i) a recombinant nucleic acid molecule encoding a glucosamine-6-phosphate acetyltransferase that <u>is at least 95% identical</u> to has the amino acid sequence of SEQ ID NO:30; and
  - ii) a recombinant nucleic acid molecule encoding a glucosamine-6-phosphate synthase that <u>is at least 95% identical to has</u> the amino acid sequence of SEQ ID NO:6; and
  - b) collecting a product produced from the step of culturing which is selected from the group consisting of glucosamine-6-phosphate, glucosamine, glucosamine-1-phosphate, N-acetylglucosamine-1-phosphate, N-acetylglucosamine-6-phosphate[[,]] and N-acetylglucosamine.
- 223. (Previously Presented) The method of Claim 229, wherein the *E. coli* further comprises a partial or complete deletion of *nagA*, *nagB*, and *nagE*.
- 224. (Previously Presented) The method of Claim 229, wherein the *E. coli* further comprises a partial or complete deletion of *manXYZ*.
- 225. (Previously Presented) The method of Claim 229, wherein the recombinant nucleic acid molecules of (a)(i) and (a)(ii) are inducible by lactose or

galactose.

- 226. (Previously Presented) The method of Claim 229, wherein the step of culturing is performed in a fermentation medium comprising glucose and fructose.
  - 227. (Canceled).
- 228. (Currently Amended) The method of claim 219, wherein the microorganism bacterium or yeast further comprises a partial or complete deletion of an endogenous gene encoding a phosphofructokinase.
- 229. (Previously Presented) The method of claim 222, wherein the *E. coli* further comprises a partial or complete deletion of *pfkA*.
- 230. (Currently Amended) The method of Claim 228, wherein the microorganism bacterium or yeast further comprises a partial or complete deletion of endogenous genes encoding N-acetylglucosamine-6-phosphate deacetylase, glucosamine-6-phosphate deaminase, and N-acetyl-glucosamine-specific enzyme II<sup>Nag</sup>.
- 231. (Currently Amended) The method of Claim 228, wherein the microorganism bacterium or yeast further comprises a partial or complete deletion of an endogenous gene encoding mannose transporter EIIM,P/III<sup>Man</sup>.
- 232. (Previously Presented) The method of Claim 228, wherein the recombinant nucleic acid molecules of (a)(i) and (a)(ii) are inducible by lactose or galactose.
- 233. (Previously Presented) The method of Claim 228, wherein the step of culturing is performed in a fermentation medium comprising glucose and fructose.

- 234. (Previously Presented) The method of Claim 219, wherein the glucosamine-6-phosphate synthase has an amino acid sequence that is at least 95% identical to SEQ ID NO:4.
- 235. (Previously Presented) The method of Claim 219, wherein the glucosamine-6-phosphate synthase has an amino acid sequence that is at least 95% identical to SEQ ID NO:6.
- 236. (Previously Presented) The method of Claim 219, wherein the glucosamine-6-phosphate synthase has an amino acid sequence that is at least 95% identical to SEQ ID NO:8.
- 237. (Previously Presented) The method of Claim 219, wherein the glucosamine-6-phosphate synthase has an amino acid sequence that is at least 95% identical to SEQ ID NO:10.
- 238. (Previously Presented) The method of Claim 219, wherein the glucosamine-6-phosphate synthase has an amino acid sequence that is at least 95% identical to SEQ ID NO:12.
- 239. (Previously Presented) The method of Claim 219, wherein the glucosamine-6-phosphate synthase has an amino acid sequence that is at least 95% identical to SEQ ID NO:14.

240-242 (Canceled).

243. (New) The method of Claim 219, wherein the glucosamine-6-phosphate synthase has an amino acid sequence that is at least 95% identical to SEQ ID NO:16.

- 244. (New) The method of Claim 219, wherein the glucosamine-6-phosphate synthase has an amino acid sequence that is at least 95% identical to SEQ ID NO:18.
- 245. (New) The method of Claim 219, wherein the glucosamine-6-phosphate synthase has an amino acid sequence that is at least 95% identical to SEQ ID NO:20.
- 246. (New) The method of Claim 219, wherein the microorganism is a bacterium or yeast.
- 247. (New) The method of Claim 219, wherein the microorganism is a bacterium.
  - 248. (New) The method of Claim 219, wherein the microorganism is a yeast.
- 249. (New) The method of Claim 222, wherein the glucosamine-6-phosphate acetyltransferase has the amino acid sequence of SEQ ID NO:30.
- 250. (New) The method of Claim 222, wherein the glucosamine-6-phosphate synthase has the amino acid sequence of SEQ ID NO:6.
  - 251. (New) The method of Claim 1, wherein the microorganism is a yeast.
- 252. (New) The method of claim 1, further comprising the step of contacting the fermentation medium with at least one ion exchange resin.